

N.G. Gemejiyeva^{1*}, D.Sh. Abdildanov², G.M. Kudabayeva³, P.V. Vesselova⁴,
N.V. Nelina⁵, N.V. Friesen⁶

^{1,3,4,5}RSE on REM “Institute of Botany and Phytointroduction” FWC MENR RK, Almaty, Kazakhstan;

²Al-Farabi Kazakh National University, Almaty, Kazakhstan;

⁶Botanical Garden of the University of Osnabrück, Germany

*Corresponding author: ngemed58@mail.ru

Species of the genus *Allium* L. of the western part of the Kyrgyz Alatau ridge: the current status and prospects of studying

The results of the assessment of the current state and prospects for the study of species of the genus *Allium* L. in the western part of the Kyrgyz Alatau ridge are presented. It was found that in the herbarium fund (AA) the most part (57 %) of specimen collections of *Allium* L. species falls on the periods from 1947 to 1960, further on descending 24 % of collections — in 1984–1985. The share of scattered collections is 12 %, the share of modern collections is insignificant (7 %). The preliminary list of studied species of *Allium* Kyrgyz Alatau is represented by 27 species. In 5 gorges of the western part of the Kyrgyz Alatau ridge, locations, phytocenotic and altitudinal distribution of 7 species were revealed. Commercial thickets are formed by *A. atosanguineum* (Kaskasu gorge) and *A. longicuspis* (Sugaty gorge). The identified species occur as part of herbaceous-grass, herbaceous-shrubby-cereal grass, petrophytic-herb-grass-shrub, petrophytic-tree-shrub communities and grow in the altitude range from 1200 to 3390 m above sea level. *Allium caeruleum*, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. longicuspis*, *A. margaritae* are found in the low and middle mountains from 1170 to 1660 metres above sea level. The high-altitude species is *A. atosanguineum*, occurring in the altitude range of 2520–3390 m above sea level. Maps of the locations and potential harvesting sites of the identified *Allium* species of the study region were compiled.

Keywords: genus *Allium* L., species diversity, Kyrgyz Alatau ridge.

Introduction

The Global Strategy for Plant Conservation [1] pays special attention to solving the issues of regional research on the inventory of species composition, identifying places of concentration of rare, endemic, relict, and economically valuable species, as well as measures to preserve botanical diversity (specific territories).

Among the economically valuable plants of Kazakhstan, wild onions are characterized by significant species diversity, wide distribution and resource potential.

At present, the genus *Allium* L. (Amaryllidaceae J.St. -Hil.) has about 1000 species [2]. About one third of onions are identified in the mountains of the Tien Shan, which is the world's largest center of onion diversity. *Allium* L. is listed in 16 major genera and is represented by 56 Tien-Shan endemics [3], of which from 9 to 21 endemic species are indicated for the territory of Kazakhstan [4, 5].

Current literature data indicate that *Allium* L. species are characterized by pronounced antioxidant properties and are promising for use in the treatment of cancer and cardiovascular diseases [6–16].

Predation of wild onions because of their ornamental, food, melliferous and medicinal properties and economic activities, including intensive grazing within their range have led to a decrease in the number of onions. The Red Data Book of Kazakhstan includes 12 species [17].

The results of a previously published review on the degree of study of 134 Kazakhstani onion species, about half of which were tested in culture, indicate the prospect of their comprehensive study in theoretical and practical terms [18].

The Kyrgyz Alatau ridge is characterized by a transitional composition of flora and a special set of species, which is associated with its location at the junction of large mountain systems of the Tien Shan. An integrated approach to their study will provide new fundamental knowledge as a scientific basis for sustainable use, conservation of plant resources and forecasting of environmental protection activities in the region.

The objects of our research are species of *Allium* L. of the Kyrgyz Alatau ridge, located within Kazakhstan.

The aim of the present research: assessment of species composition, distribution and current state of natural populations of onions of the western part of the Kyrgyz Alatau ridge for their conservation and sustainable use.

Experimental

Herbarium collections (Herbarium AA) and natural populations of onions growing in the western part of the Kyrgyz Alatau ridge served as the material for research.

The research area — the Kyrgyz Alatau ridge is located within the Zhambyl, T. Ryskulov and Merken administrative districts of Zhambyl region, characterized by a variety of relief, soil and vegetation cover, represented from high-altitude subalpine and alpine vegetation to psammophilic vegetation of Moyinkum sands.

The Kyrgyz ridge is located within the Kyrgyz Mountain Subprovince, which is part of the Dzungaro-North Tianshan Mountain Province [19].

Common methods of botanical and cartographic research were used in the work.

Classical geobotanical methods [20–22] were used in the expedition studies. Determination of plant species was carried out according to floristic summaries: “Flora of Kazakhstan” [23–31], “Illustrated identifier...” [32–33]. The taxonomy of species is agreed with the Internet resources: Plantarium [34] and Plants of the World Online (POWO) [35], but the authors adhere to their own point of view on the species affiliation of some taxa. The distribution of the studied species by nature of use and pharmacological action was carried out according to the multi-volume summary: “Plant Resources...” [36–38], “Wild Usable Plants of Russia” [39], “Annotated list of medicinal plants of Kazakhstan” [40].

The digital geographic base of 1:2500000 scale of FGBU “VSEGEI” was used as a cartographic basis for mapping, and updated by the authors’ own efforts. Coordinates of the area, where the locations of bows were identified, were recorded in the field using GNSS-navigator “Garmin”. To create maps of locations of fishery arrays, the coordinates obtained with the help of the navigator were plotted on the map base and processed in ESRI ArcGIS Desktop 10.8 software.

Results and Discussion

To identify species diversity and compile a preliminary list of onions of the studied region, including rare and endemic representatives, as well as their potential locations, we screened available herbarium material, literature and field data, and Internet resources. Field studies were carried out, primary material was collected and locations of identified *Allium* L. species of the study area were mapped.

As a result of screening of herbarium materials of representatives of the genus *Allium* L. of the Kyrgyz Alatau in the herbarium collection (AA), 82 specimens were reviewed, including 3 specimens of the rare species *Allium suworowii* Regel [17] (Fig. 1).

Analysis of the herbarium specimens viewed showed that the majority of herbarium specimens (19 or 23 %) represented the species *A. barsczewskii* Lipsky. For other species, the number of herbarium specimens varies from 2 to 7: *A. caesium* Schrenk, *A. fetisowii* Regel — 7 (8.5 %), *A. caeruleum* Pall. — 5 (6.1 %), *A. trachyscordum* Vved., *A. karataviense* Regel, *A. hymenorhizum* Ledeb., *A. atosanguineum* Kar. et Kir., *A. platyspathum* Schrenk, *A. caricifolium* Kar. & Kir. (syn. *A. pallasii* Murray), *A. oreoprasum* Schrenk — 3 (3.6 %), *A. parvulum*, *A. oreophilum*, *A. oreoprasoides* Vved. — 2 (2.4 %). Eight species (1 (1.2 %) herbarium specimen each) are represented in the lowest number: *A. obliquum* L., *A. turkestanicum* Regel, *A. semenovii* Regel, *A. longicuspis* Regel, *A. kokanicum* Regel, *A. schoenoprasum* L., *A. longiradiatum* (Regel) Vved. It was revealed that the most part (57 %) of collections of species of the genus *Allium* L. falls on the periods from 1947 to 1960, further in descending order 24 % of collections — on 1984–1985. The share of collections of different years is 12 %, modern collections — insignificant (7 %).

To compile a preliminary list of species of the genus *Allium* L. in the Kazakh part of the Kyrgyz Alatau ridge, the analysis of available data on onions occurring in the Kyrgyz Alatau was continued. Comparison of published data on the distribution of species of the genus *Allium* L. according to floristic summaries [24, 32, 41]; Internet resources [34] and herbarium data (AA) showed that for 7 species (*A. caeruleum* Pall. *A. obliquum* L., *A. oreoprasoides* Vved., *A. turkestanicum* Regel, *A. longiradiatum* (Regel) Vved., *A. suworowii* Regel, *A. caricifolium* Kar. & Kir. (syn. *A. pallasii* Murr.) found new localities on the territory of the Kyrgyz Alatau that not indicated in the literature sources. At the same time, the onion species of the Kyrgyz ridge *A. altissimum* Regel and *A. leptomorphum* Vved. indicated in the literature sources are absent in the herbarium collection (AA) (Table 1).



Figure 1. Herbarium specimens of some species of the genus *Allium* collected in the Kyrgyz Alatau

According to the “Flora of Kazakhstan” [24] there are 12 species of onions in the Kyrgyz Alatau; in the “Illustrated Plant Identifier of Kazakhstan” [32] there are 15 species; in the “Plant Identifier of Central Asia” [32] there are 15 species of onions; in the “Plant Identifier of Central Asia” [41] — 17 species [41]. The localities of *A. margaritae* B. Fedtsch. in the Kyrgyz Alatau are also absent in published sources (Table 1).

In the Red data book of Zhambyl region [42], out of 5 specified representatives of the genus *Allium* L.

(*A. galanthum* Kar. & Kir., *A. longicuspis*, *A. kasteki*, *A. karataviense*, *A. turtschicum*), only one rare species *A. karataviense* is listed for the Almalysai gorge of the Kyrgyz Alatau.

Based on screening of available herbarium material and analysis of published data, a preliminary list of species of *Allium* L. Kyrgyz Alatau is represented by 27 species was compiled: *Allium altissimum* Regel, *A. atosanguineum* Schrenk, *A. barsczewskii* Lipsky, *A. caeruleum* Pall., *A. caesium* Schrenk, *A. caricifolium* Kar. & Kir. (syn. *A. pallasii* Murray), *A. dasyphyllum* Vved., *A. fetisowii* Regel, *A. hymenorhizum* Ledeb., *A. karataviense* Regel, *A. kokanicum* Regel, *A. leptomorphum* Vved., *A. longicuspis* Regel, *A. longiradiatum* (Regel) Vved., *A. margaritae* B. Fedtsch., *A. oreophilum* C.A. Mey., *A. oreoprasum* Schrenk, *A. oreoprasoides* Vved., *A. obliquum* L., *A. parvulum* Vved., *A. platyspathum* Schrenk, *A. polyphyllum* Kar. & Kir., *A. schoenoprasum* L., *A. semenovii* Regel, *A. suworowii* Regel, *A. trachyscordum* Vved., *A. turkestanicum* Regel, of which 1 endemic (*A. oreoprasoides*), 2 sub-endemics (*A. margaritae*, *A. trachyscordum*), which are also distributed in Kyrgyzstan, and 1 rare species (*A. suworowii*).

Table 1

Comparative data on the presence of species of the genus *Allium* L. in the flora of the Kyrgyz Alatau according to floristic summaries, herbarium (AA) and Internet resources

Name of the species by FK	Species name adopted by POWO [35]	FK [24]	IIPK [32]	PICA [41]	Herbarium (AA)	Plantarium [34]
<i>Allium altissimum</i> Regel	<i>Allium altissimum</i> Regel	+	+	+	—	—
<i>A. atrosanguineum</i> Kar. & Kir.	<i>A. atrosanguineum</i> Schrenk	—	+	+	+	—
<i>A. barsczewskii</i> Lipsky	<i>A. barsczewskii</i> Lipsky	+	+	+	+	—
<i>A. caeruleum</i> Pall.	<i>A. caeruleum</i> Pall.	—	+	+	+	—
<i>A. caesium</i> Schrenk	<i>A. caesium</i> Schrenk	—	+	+	+	+
<i>A. caricifolium</i> Kar. & Kir.	syn. <i>A. pallasii</i> Murray	—	—	—	+	—
<i>A. dasyphyllum</i> Vved.	<i>A. dasyphyllum</i> Vved.	—	—	+	+	+
<i>A. fetisowii</i> Regel	<i>A. fetisowii</i> Regel	+	+	+	+	+
<i>A. hymenorrhizum</i> Ledeb.	<i>A. hymenorrhizum</i> Ledeb.	+	+	+	+	—
<i>A. karataviense</i> Regel	<i>A. karataviense</i> Regel	—	—	—	+	+
<i>A. kokanicum</i> Regel	<i>A. kokanicum</i> Regel	+	+	+	—	—
<i>A. leptomorphum</i> Vved.	<i>A. leptomorphum</i> Vved.	—	—	+	—	—
<i>A. longicuspis</i> Regel	<i>A. sativum</i> L.	—	—	—	+	—
<i>A. longiradiatum</i> (Regel) Vved.	<i>A. longiradiatum</i> (Regel) Vved.	—	—	—	+	—
<i>A. margaritae</i> B. Fedtsch.	<i>A. margaritae</i> B. Fedtsch.	—	—	—	+	+
<i>A. oreophilum</i> C.A. Mey.	<i>A. oreophilum</i> C.A. Mey.	+	+	+	—	—
<i>A. oreoprasum</i> Schrenk	<i>A. oreoprasum</i> Schrenk	+	+	+	+	—
<i>A. oreoprasoides</i> Vved.	<i>A. oreoprasoides</i> Vved.	—	—	—	+	—
<i>A. obliquum</i> L.	<i>A. obliquum</i> L.	—	—	—	+	—
<i>A. parvulum</i> Vved.	<i>A. parvulum</i> Vved.	+	+	+	+	+
<i>A. platyspathum</i> Schrenk	<i>A. platyspathum</i> Schrenk ex Fisch. & C.A. Mey.	+	+	+	+	—
<i>A. polyphyllum</i> Kar. & Kir.	<i>A. carolinianum</i> Redouté	+	+	+	+	—
<i>A. schoenoprasum</i> L.	<i>A. schoenoprasum</i> L.	—	—	—	+	—
<i>A. semenovii</i> Regel	<i>A. semenovii</i> Regel	+	+	+	+	—
<i>A. suworowii</i> Regel	<i>A. suworowii</i> Regel	—			+	
<i>A. trachyscordum</i> Vved.	<i>Allium trachyscordum</i> Vved.	+	+	+	+	+
<i>A. turkestanicum</i> Regel	<i>A. turkestanicum</i> Regel	—	—	—	+	—
	Total species	12	15	17	23	7

Note – POWO — Plants of the World Online; FK — Flora of Kazakhstan; IIPK — Illustrated identifier of plants of Kazakhstan; PICA — Plant Identifier of Central Asia

Since the territory of the Kyrgyz Alatau ridge is transboundary, it is possible to expand the preliminary list with other species of onions growing in Kyrgyzstan. Of the 87 species of the genus *Allium* L. identified in Kyrgyzstan, 24 grow in both Kyrgyzstan and Kazakhstan [43–46].

The ridge of application of onions in the Kazakh part of the Kyrgyz Alatau is very diverse. Six species are used as medicinal (*Allium altissimum*, *A. karataviense*, *A. longicuspis*, *A. obliquum*, *A. schoenoprasum*, *A. suworowii*). Ornamental properties are characterized by 16 species, food — 9; honey-bearing — 5 species, fodder, technical and poisonous — 1 species each. 21 species have been tested in culture.

In August 2024, 5 gorges of the western part of the Kyrgyz Alatau (Syugaty, Kaskasu, Karakystak, Kursai, Ulken Almalysai) within Zhambyl, Merken and T. Ryskulov administrative districts of Zhambyl region were surveyed. The points of occurrence, altitudinal confinement, distribution of 7 species of the genus *Allium* L. (*A. atrosanguineum*, *A. caeruleum*, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. longicuspis*, *A. margaritae*).

Field surveys in 2024 showed that the species identified in the gorges of the western part of the Kyrgyz Alatau ridge occurred in the altitude range from 1200 to 3390 m a.s.l. Locations of six species (*Allium*

caeruleum, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. margaritae*, *A. longicuspis*) were recorded in the low and middle mountains in the altitude range from 1170 to 1660 m a.s.l., and the high-mountain species *A. atosanguineum* grew in the upper reaches of the Karakystak and Kaskasu rivers in the altitude range of 2520–3390 m a.s.l.

It should be noted that from 2 to 4 species of bows were found in the surveyed gorges: Syugates (*A. caeruleum*, *A. caesium*, *A. karataviense*, *A. longicuspis*), Ulken Almalysai (*A. caeruleum*, *A. karataviense*, *A. margaritae*), Karakystak (*A. atosanguineum*, *A. leptomorphum*), Kursai (*A. caeruleum*, *A. karataviense*), with the exception of the Kaskasu gorge (*A. atosanguineum*). At the same time, the locations of *A. caeruleum* and *A. karataviense* were recorded in three gorges (Kursai, Syugaty, Ulken Almalysai) in the western part of the Kyrgyz Alatau in the altitude ridge 1200–1440 m above sea level (Table 2).

The identified onions were often found as part of herbaceous-grass, herbaceous-shrubby-cereal grass, petrophytic-herb-grass-shrub and petrophytic-tree-shrub communities. Over 32 species of vascular plants were also observed in the above communities, along with onions: *Acer semenovii* Regel & Herder, *Achillea millefolium* L., *Althaea officinalis* L., *Artemisia dracunculus* L., *Artemisia serotina* Bunge, *Bassia prostrata* (L.) Beck, *Bromus japonicus* Thunb., *Cerasus tianshanica* Pojark., *Convolvulus arvensis* L., *Ephedra intermedia* Schrenk & C.A. Mey., *Euphorbia jaxartica* (Prokh.) Krylov, *Ferula diversivittata* Regel & Schmalh., *Galium aparine* L., *Galium verum* L., *Haplophyllum acutifolium* (DC.) G. Don, *Hypericum perforatum* L., *Hypericum scabrum* L., *Lomelosia songarica* (Schrenk) Soják, species of the genus *Lonicera* L., *Megacarpaea orbiculata* B. Fedtsch., *Mentha asiatica* Boriss., *Onosma irritans* Popov, *Origanum vulgare* L., *Poa bulbosa* L., *Rhaponticoides ruthenica* (Lam.) M.V. Agab. & Greuter, *Rheum tataricum* L. f., *Rosa persica* Michx. ex Juss., species of the genus *Rosa* L., *Sisymbrium altissimum* L., *Sisymbrium loeselii* L., *Sonchus palustris* L., *Spiraea hypericifolia* L., *Stipa caucasica* Schmalh., *Tulipa greigii* Regel etc. from 19 families (Fig. 2).

Maps of locations of studied onions in the region were created based on literature, herbarium and Plantarium data (Fig. 3), results of expedition studies (Fig. 4), indicating places of potential harvesting of *Allium* L. species identified in 2024.

Table 2

Distribution and altitudinal habitat of identified species of the genus *Allium* L. in the western part of the Kyrgyz Alatau ridge according to field surveys in 2024

Species	Coordinates	Elevation above sea level	Location, date, year
<i>Allium atosanguineum</i> Kar. & Kir.	N 42°32'37" E 73°12'12"	3390 m	Zhambyl region, Merken district, Kyrgyz Alatau ridge, Kaskasu village, south-eastern slope. Kaskasu, south-eastern slope. 28.08.2024.
	N 42°39'53" E 72°47'12"	2520 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, Karakystak gorge, southern slope. 28.08.2024.
<i>Allium caeruleum</i> Pall.	N 42°53'54" E 71°44'20"	1440 m	Zhambyl region, Zhambyl district, Kyrgyz Alatau ridge, gorge Kursai. 26.08.2024.
	N 42°51'49" E 71°50'56"	1290 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, gorge Syugaty, along the road, on an earthen slope. 27.08.2024.
	N 42°52'9" E 71°49'41"	1280 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, Syugaty village, south-eastern slope, gorge Syugaty, south-eastern slope. 27.08.2024.
	N 42°52'12" E 71°49'40"	1270 m	Zhambyl region, T. Ryskulov district, Kirghiz Alatau ridge, gorge Syugaty, lower part of stony-rubble slope of southern exposure. 27.08.2024.
	N 42°53'56" E 71°42'43"	1200 m	Zhambyl region, Zhambyl district, Kyrgyz Alatau ridge, gorge Ulken Almalysay, southern slope. 26.08.2024.
<i>Allium caesium</i> Schrenk	N 42°51'48" E 71°51'1"	1320 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, gorge Syugaty, stony-rubble slope of southern exposure. 27.08.2024.

Continuation of Table 2

Species	Coordinates	Elevation above sea level	Location, date, year
<i>Allium karataviense</i> Regel	N 42°53'54" E 71°44'20"	1440 m	Zhambyl region, Zhambyl district, Kyrgyz Alatau ridge, gorge Kursai. 26.08.2024.
	N 42°52'12" E 71°49'40"	1270 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, gorge Syugaty, lower part of stony-rubble slope of southern exposure. 27.08.2024.
	N 42°53'56" E 71°42'43"	1200 m	Zhambyl region, Zhambyl district, Kyrgyz Alatau ridge, gorge Ulken Almalysay, southern slope. 26.08.2024.
<i>Allium leptomorphum</i> Vved.	N42°37'33" E72°53'29"	1660 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, gorge Karakystak, southern slope. 28.08.2024.
<i>Allium longicuspis</i> Regel	N 42°52'17" E 71°50'13"	1170 m	Zhambyl region, T. Ryskulov district, Kyrgyz Alatau ridge, gorge Syugaty. 26.08.2024.
<i>Allium margaritae</i> B. Fedtsch.	N42°53'56" E71°42'43"	1200 m	Zhambyl region, Zhambyl district, Kyrgyz Alatau ridge, gorge Ulken Almalysay, southern slope. 26.08.2024.




Figure 2. Distribution of associated plant species by families




Figure 3. Map* of the locations of the studied species of the genus *Allium* L. in the western part of the Kyrgyz Alatau according to literature, herbarium (AA) and Plantarium data (* — author of the map Rybakov I.A.)




Figure 4. Map* of locations and places of potential harvesting of identified species of the genus *Allium* L. in the western part of the Kyrgyz Alatau in 2024 (* — author of the map Rybakov I.A.)

Conclusion

Modern assessment of species composition, distribution and state of natural populations of bows of the western part of the Kyrgyz Alatau Ridge showed that the majority (57 %) of collections of species of *Allium* L. in the herbarium collection (AA) are from 1947 to 1960, followed by 24 % of collections from 1984–1985. The proportion of collections from different years is 12 % and modern collections are negligible (7 %). The preliminary list of species of the genus *Allium* L. in the Kazakh part of the Kyrgyz Alatau ridge is represented by 27 species, including 1 endemic, 2 subendemic and 1 rare species. It is possible to expand the preliminary list with other species of onions growing in the transboundary territory with Kyrgyzstan.

In 5 gorges of the western part of the Kirghiz Alatau ridge, the locations, phytocoenotic and altitudinal distribution of 7 species were revealed, of which *A. atosanguineum* (Kaskasu gorge) and *A. longicuspis* (Syugaty gorge) form commercial thickets. The studied bows were found in the altitude ridge from 1200 to 3390 m a.s.l. They were found as part of various plant communities, including petrophytic communities. In the lowlands and midlands from 1170 to 1660 m a.s.l.: *Allium caeruleum*, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. longicuspis*, *A. margaritae*.

The high-altitude species is *A. atosanguineum*, occurring in the altitude range of 2520–3390 m above sea level. Locations and potential harvesting sites of the studied species of the genus *Allium* L. of the western part of the Kyrgyz Alatau are presented on cartographic models. Floristic and resource studies of onions continue.

Funding

This research is funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant AP23484591 “Comprehensive research (floristic, molecular genetic and resource) on the *Allium* L. genus of the Kyrgyz Alatau cross-border ridge — a scientific foundation for the management of the region’s natural resources” (2024–2026)).

Acknowledgment

We express our gratitude to I.A. Rybakov, cartographer, senior researcher of the laboratory of plant resources, for the created cartographic models.

References

- 1 Глобальная стратегия сохранения растений: 2011–2020 (Convention on Biological Diversity, 2012. Global Strategy for Plant Conservation: 2011–2020). — 2002. — Botanic Gardens Conservation International, Richmond, UK.
- 2 Govaerts R. World checklist of Amaryllidaceae / R. Govaerts, S. Kington, N. Friesen, R. Fritsch, D.A. Snijman, R. Marcucci, P.A. Silverstone-Sopkin, S. Brullo. — 2024. — [Electronic resource]. — Access mode: <http://apps.kew.org>
- 3 Tojibaev K.S. An Annotated Checklist of Endemic Vascular Plants of the Tian-Shan Mountains in Central Asian Countries / K.S. Tojibaev, C.G. Jang, G.A. Lazkov, K.S. Chang, G.T. Sitpayeva, N. Safarov, S.H. Oh // Phytotaxa. — 2020. — Vol. 464, No. 2. P. — 117–158. DOI: 10.11646/phytotaxa.464.2.1
- 4 Об утверждении перечня эндемичных и реликтовых растений. Приказ Министра экологии и природных ресурсов Республики Казахстан от 7 марта 2023 года № 78. — [Электронный ресурс]. — Режим доступа: <https://adilet.zan.kz/rus/docs/V2300032050/links>
- 5 Kubentayev S.A. Revised checklist of endemic vascular plants of Kazakhstan / S.A. Kubentayev, D.T. Alibekov, Y.V. Perezhogin, G.A. Lazkov, A.N. Kupriyanov, A.L. Ebel, K. Izbastina, O. Borodulina, B.B. Kubentayeva // PhytoKeys. — 2024. — Vol. 238. — P. 241–279. <https://doi.org/10.3897/phytokeys.238.114475>.
- 6 Steiner D. Comparative study on *Allium schoenoprasum* L. cultivated plant and *Allium schoenoprasum* tissue culture organs antioxidant status / D. Steiner, B.M. Popovic, D. Calic-Dragosavac, D. Malencic, and S. Zdravcovic-Korac // Phytother. res. — 2011. — Vol. 25. — P. 1618–1622.
- 7 Ширшова Т.И. Содержание селена в культурных и дикорастущих луках из флоры Республики Коми / Т.И. Ширшова, И.В. Бешлей, Н.В. Матистов, Н.А. Голубкина // Растительные ресурсы. — 2011. — Т. 47. — Вып. 1. — С. 112–118.
- 8 Ширшова Т.И. Изучение химического состава листьев *A. schoenoprasum* L. и ингибирующего действия их экстрактов на опухолевый рост у мышей / Т.И. Ширшова, И.В. Бешлей, В.П. Дерягина, Н.И. Рыжова, Н.В. Матистов // Хим.-фарм. журн. — 2012. — Т. 46. — № 11. — С. 40–43. DOI: <https://doi.org/10.30906/0023-1134-2012-46-11-30-33>.
- 9 Матистов Н.В. Микронутриенты дикорастущих и культивируемых видов рода *Allium* (*A. angulosum*, *A. strictum*, *A. schoenoprasum*) на европейском Северо-Востоке России: автореф. дис. ... канд. биол. наук / Н.В. Матистов. — Новосибирск, 2013. — 22 с.

- 10 Jang J.E. Flower morphology of *Allium* (Amaryllidaceae) and its systematic significance / J.E. Jang, S. Baasanmunkh, N. Nyamgerel, S.Y. Oh, J.H. Song, Z. Yusupov, K. Tojibaev, H.J. Choi // *Plant Diversity*. — 2024. — Vol. 46. — No. 1. — P. 3–27 DOI: 10.1016/j.pld.2023.06.009.
- 11 Ivanova M.I. Assessment of phytochemicals in *Allium* species: a systematic review / M.I. Ivanova, A.A. Baikov, E.M. Gins, V.K. Gins, A.I. Kashleva, M.S. Gins, N.V. Smurova // *Sabrao Journal of Breeding & Genetics*. — 2024. — Vol. 56. — No. 3. DOI: 10.54910/sabrao2024.56.3.13.
- 12 Teotia D. Pharmacophylogeny of genus *Allium* L. / D. Teotia, A. Agrawal, H. Goyal, P. Jain, V. Singh, Y. Verma, K. Parveen, N.A. Bukhari, A. Chandra, V. Malik // *Journal of King Saud University-Science*. — 2024. — Vol. 36. — No. 8. — P. 103330 DOI: 10.1016/j.jksus.2024.103330.
- 13 Feknous N. Updated Insights on the Antimicrobial Activities of *Allium* Genus (A Review) / N. Feknous, M. Boumendjel, F.Z. Leblab // *Russian Journal of Bioorganic Chemistry*. — 2024. — Vol. 50. — No. 3. — P. 806–823. DOI: 10.1134/S1068162024030257.
- 14 Iwar K. *Allium* as Potential Antioxidants and Anticancer Agents / K. Iwar, K. Ochar, Y.A. Seo, B.K. Ha, S.H. Kim // *International Journal of Molecular Sciences*. — 2024. — Vol. 25. — No. 15. — P. 8079 DOI: 10.3390/ijms25158079.
- 15 Friesen N. Dated phylogeny, phylogeography, and classification of *Allium* subgenus *Amerallium* (Amaryllidaceae) from the Old World, based on six DNA fragments / N. Friesen, T. Herden, M. Leweke, L. Grützmacher, O. Fragman-Sapir, H. Hurka, F.R. Blattner, R.M. Fritsch // *TAXON*. — 2024. — Vol. 73. — No. 4. — P. 971–991. DOI: <https://doi.org/10.1002/tax.13217>.
- 16 Vatandaşlar Ö. Comparative Seed Morphology of Eight *Allium* L. Species from Sections Codonoprasum and Scorodon / Ö. Vatandaşlar, M. Koçyiğit // *Turkish Journal of Bioscience and Collections*. — 2024. — Vol. 8. — No. 2. — P. 41–48 DOI: <https://doi.org/10.26650/tjbc.1525986>.
- 17 Красная книга Казахстана / гл. ред. И.О. Байтулин, отв. ред. Г.Т. Ситпаева. — изд. 2-е, исправл. и дополн. — Астана: ТОО «ArtPrintXX1», 2014. — Т. 2. Ч. 1. Растения. — 452 с.
- 18 Гемеджиева Н.Г. Обзор современного состояния и перспективы изучения казахстанских видов рода *Allium* L. / Н.Г. Гемеджиева, А.М. Токенова, Н.В. Фризен // *Проблемы ботаники Южной Сибири и Монголии*, 2021. — Т. 20, № 1. — С. 97–101. DOI: <https://doi.org/10.14258/pbssm.2021020>.
- 19 Рачковская Е.И. Принципы и основные единицы районирования / Е.И. Рачковская, И.Н. Сафронова, Е.А. Волкова; под редакцией Е.И. Рачковской, Е.А. Волковой, В.Н. Храмцова // *Ботаническая география Казахстана и Средней Азии (в пределах пустынной области)*. — СПб., 2003. — С. 172–195.
- 20 Корчагин А.А. Видовой (флористический) состав растительных сообществ и методы его изучения / А.А. Корчагин // *Полевая геоботаника*. — М.–Л., 1964. — Т. 3. — С. 39–60.
- 21 Понятовская В.М. Учет обилия и особенности размещения видов в естественных растительных сообществах / В.М. Понятовская // *Полевая геоботаника*. — М.–Л., 1964. — Т. 3. — С. 209–237.
- 22 Быков Б.А. Геоботаника / Б.А. Быков. — Алма-Ата: Наука, 1978. — 288 с.
- 23 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1956. — Т. 1. — 354 с.
- 24 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1958. — Т. 2. — 292 с.
- 25 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1960. — Т. 3. — 460 с.
- 26 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1961. — Т. 4. — 548 с.
- 27 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1961. — Т. 5. — 515 с.
- 28 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1963. — Т. 6. — 465 с.
- 29 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1964. — Т. 7. — 497 с.
- 30 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1965. — Т. 8. — 447 с.
- 31 Флора Казахстана: [в 9-и т.] / ред. Н.В. Павлов. — Алма-Ата: Изд-во АН КазССР, 1966. — Т. 9. — 640 с.
- 32 Иллюстрированный определитель растений Казахстана [в 2-х т.] / ред. В.П. Голоскоков. — Алма-Ата: Изд-во «Наука» Казахской ССР, 1969. — Т. 1. — 644 с.
- 33 Иллюстрированный определитель растений Казахстана [в 2-х т.] / ред. В.П. Голоскоков. — Алма-Ата: Изд-во «Наука» Казахской ССР, 1972. — Т. 2. — 572 с.
- 34 Плантариум: открытый онлайн атлас-определитель растений и лишайников России и сопредельных стран. — 2007–2024. — [Электронный ресурс]. — Режим доступа: <http://www.plantarium.ru/>
- 35 Plants of the World Online. — POWO, 2024. — [Electronic resource]. — Access mode: <https://powo.science.kew.org>
- 36 Растительные ресурсы России и сопредельных государств: Цветковые растения, их химический состав, использование. Сем. Butomaceae — Турпасеae / отв. ред. П.Д. Соколов. — СПб.: Наука, 1994. — 271 с.
- 37 Растительные ресурсы России и сопредельных государств. Цветковые растения, их химический состав, использование / отв. ред. А.Л. Буданцев. Ч. 1: Семейства Lysorodiaceae — Ephedraceae; Ч. 2: Дополнения к 1–7 томам. — СПб.: Мир и семья–95, 1996. — 571 с.
- 38 Растительные ресурсы России. Дикорастущие цветковые растения, их компонентный состав и биологическая активность. — Том 1. Семейства Magnoliaceae — Juglandaceae, Ulmaceae, Moraceae, Cannabaceae, Urticaceae / отв. ред. А.Л. Буданцев. — СПб.; М.: Товарищество научных изданий КМК, 2008. — 421 с.
- 39 Дикорастущие полезные растения России / отв. ред: А.Л. Буданцев, Е.Е. Лесиовская. — СПб.: Издательство СПХФА, 2001. — С. 400.

- 40 Аннотированный список лекарственных растений Казахстана: Справочное издание / Л.М. Грудзинская, Н.Г. Гемеджиева, Н.В. Нелина, Ж.Ж. Каржаубекова. — Алматы, 2014. — 200 с.
- 41 Введенский А.И. Род *Allium* L. / А.И. Введенский; ред. С.С. Ковалевская // Определитель растений Средней Азии. Критический конспект флоры. — Ташкент: Изд-во ФАН Узбекской ССР, 1971. — Т. 2. — С. 39–89.
- 42 Красная книга Жамбылской области / Н.К. Аралбай, С.К. Мухтубаева, А.А. Шорманова. — Алматы, 2007. — 160 с.
- 43 Лазьков Г.А. Кадастр флоры Кыргызстана. Сосудистые растения / Г.А. Лазьков, Б.А. Султанова. — Бишкек, 2014. — С. 21–23.
- 44 Турдуматова Н.К. Род лук (*Allium* L.) в Кыргызстане (вопросы систематики, география и перспективы использования): автореф. дис. ... канд. биол. наук / Н.К. Турдуматова. — Бишкек, 2022. — 22 с.
- 45 Friesen N. *Allium pallasii* and *A. caricifolium* — surprisingly diverse old steppe species, showing a clear geographical barrier in the area of Lake Zaysan / N. Friesen, L. Grützner, M. Skaptsov, P. Vesselova, V. Dorofeyev, A.N. Luferov, N. Turdumatova, G. Lazkov, S.V. Smirnov, A.I. Shmakov, H. Hurka // Plants. — 2022. — Vol. 11. — No 11. — P. 1465. doi.org/10.3390/plants11111465.
- 46 Нелина Н.В. К видовому составу флоры ущелий Мерке и Каракыстак западной части Киргизского Алатау / Н.В. Нелина, Г.М. Кудабаяева, С.К. Мухтубаева, Б.К. Билибаева // Проблемы ботаники Южной Сибири и Монголии: мат. XV Междунар. научно-практ. конф. — Барнаул, 2016. — С. 392–398.

Н.Г. Гемеджиева, Д.Ш. Абдилданов, Г.М. Кудабаяева, П.В. Веселова,
Н.В. Нелина, Н.В. Фризен

Кыргыз Алатауы жотасынын батыс бөлігіндегі *Allium* L. туысынын түрлері: қазіргі жағдайы және зерттеу перспективалары

Мақалада Кыргыз Алатауы жотасынын батыс бөлігіндегі *Allium* L. туысынын түрлерін зерттеудің қазіргі жағдайы мен болашағын бағалау нәтижелері берілген. Гербарий қорында (АА) *Allium* L. туысы түрлерінің коллекцияларының көпшілігі (57%) 1947–1960 жылдар аралығына, одан кейін кему ретімен коллекциялардың 24%-ы 1984–1985 жылдарға келетіні анықталды. Жекелеген терімдердің үлесі 12% құрайды, ал қазіргі терімдердің үлесі шамалы (7%). Кыргыз Алатауында зерттелген *Allium* туысынын алдын-ала тізімі 27 түрден тұрады. Жотаның батыс бөлігіндегі 5 шатқалда 7 түрдің кездесетін орны, фитоценоздық және биіктік бойынша таралуы анықталды. Өндірістік қопаларды *A. atosanguineum* (Қасқауы шатқалы) және *A. longicuspis* (Сыуғаты шатқалы) құрайды. Анықталған түрлер көбінесе әртүрлі шөпті-дәнді, әр түрлі шөпті-бұталы-дәнді, жартас өсімдікті-әртүрлі шөпті-бұталы, жартас өсімдікті-ағашты-бұталы қауымдастықтарда кездеседі және теңіз деңгейінен 1200-ден 3390 м-ге дейінгі биіктік аралығында өседі. Теңіз деңгейінен 1170–1660 м биіктіктегі аласа және орташа таулы аймақтарда *Allium caeruleum*, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. longicuspis*, *A. margaritae* анықталды. Сонымен қатар *A. atosanguineum* биік таулы түр, ол теңіз деңгейінен 2520–3390 м биіктік аралығында кездеседі. Зерттелген аймақтан анықталған *Allium* түрлерінің кездесетін жері мен әлеуметтік жинау орындарының карталары құрастырылды.

Кілт сөздер: түрлердің әртүрлілігі, *Allium* L. туысы, алуантүрлілік, Кыргыз Алатауы жотасы.

Н.Г. Гемеджиева, Д.Ш. Абдилданов, Г.М. Кудабаяева, П.В. Веселова,
Н.В. Нелина, Н.В. Фризен

Виды рода *Allium* L. западной части хребта Киргизский Алатау: современное состояние и перспективы изучения

В данной работе приведены результаты оценки современного состояния и перспектив изучения видов рода *Allium* L. западной части хребта Киргизский Алатау. Установлено, что в гербарном фонде (АА) большая часть (57 %) коллекционных сборов видов *Allium* L. приходится на периоды с 1947 по 1960 годы, далее по убыванию 24 % сборов — на 1984–1985 годы. Доля разрозненных сборов составляет 12 %, доля современных сборов незначительна (7 %). Предварительный список изучаемых видов *Allium* в Киргизском Алатау представлен 27 видами. В 5 ущельях западной части хребта выявлены местонахождения, фитоценотическая и высотная приуроченность 7 видов. Промысловые заросли образуют *A. atosanguineum* (ущелье Каскасу) и *A. longicuspis* (ущелье Сюгаты). Выявленные виды часто встречаются в составе разнотравно-злакового, разнотравно-кустарниково-злакового, петрофитно-разнотравно-кустарникового, петрофитно-древесно-кустарникового сообществ и произрастают в интервале высот от 1200 до 3390 м над ур. м. В низкорослые и среднегорье от 1170 до 1660 м над ур. м. отмечены *Allium caeruleum*, *A. caesium*, *A. karataviense*, *A. leptomorphum*, *A. longicuspis*, *A. margaritae*.

Высокогорным видом является *A. atosanguineum*, встречающийся в интервале высот 2520–3390 м над ур. м. Составлены карты местонахождений и потенциальных мест заготовок выявленных видов *Allium* исследуемого региона.

Ключевые слова: видовое разнообразие, род *Allium* L., хребет Киргизский Алатау.

References

- 1 (2002). *Globalnaia strategiia sokhraneniia rastenii: 2011–2020 (Convention on Biological Diversity, 2012. Global Strategy for Plant Conservation: 2011–2020)*. Botanic Gardens Conservation International, Richmond, UK [in Russian].
- 2 Govaerts, R., Kington, S., Friesen, N., Fritsch, R., Snijman, D.A., Marcucci, R., Silverstone-Sopkin, P.A. & Brullo, S. (2024). World Checklist of Amaryllidaceae. Retrieved from <http://wcsp.science.kew.org>
- 3 Tojibaev, K.S., Jang, C.G., Lazkov, G.A., Chang, K.S., Sitpayeva, G.T., Safarov, N., & Oh, S.H. (2020). An annotated checklist of endemic vascular plants of the Tian-Shan Mountains in Central Asian countries. *Phytotaxa*, 464(2), 117–158. DOI: <https://doi.org/10.11646/phytotaxa.464.2.1>
- 4 Ob utverzhdenii perechnia endemichnykh i reliktovykh rastenii. Prikaz Ministra ekologii i prirodnnykh resursov Respubliki Kazakhstan ot 7 marta 2023 goda № 78. Zaregistririvan v Ministerstve yustitsii Respubliki Kazakhstan 13 marta 2023 goda № 32050 [On approval of the list of endemic and relict plants. Order of the Minister of Ecology and Natural Resources of the Republic of Kazakhstan dated 7 March 2023 No 78. Registered with the Ministry of Justice of the Republic of Kazakhstan on 13 March 2023 No. 32050]. Retrieved from <https://adilet.zan.kz/rus/docs/V2300032050/links> [in Russian].
- 5 Kubentayev, S.A., Alibekov, D.T., Perezhogin, Y.V., Lazkov, G.A., Kupriyanov, A.N., Ebel, A.L., Izbastina, K., Borodulina, O., & Kubentayeva, B.B. (2024). Revised checklist of endemic vascular plants of Kazakhstan. *PhytoKeys*, 238, 241–279. <https://doi.org/10.3897/phytokeys.238.114475>.
- 6 Štajner, D., Popović, B.M., Čalić-Drăgosavac, D., Malenčić, Đ., & Zdravković-Korać, S. (2011). Comparative study on *Allium schoenoprasum* cultivated plant and *Allium schoenoprasum* tissue culture organs antioxidant status. *Phytotherapy Research*, 25, 1618–1622. doi: 10.1002/ptr.3394.
- 7 Shirshova, T.I., Beshlej, I.V., Matistov, N.V., & Golubkina, N.A. (2011). Soderzhanie selena v kulturnykh i dikorastushchikh lukakh iz flory Respubliki Komi [Selenium content in cultivated and wild onions from the flora of the Komi Republic]. *Rastitelnye resursy — Plant Resources*, 47(1), 112–118 [in Russian].
- 8 Shirshova, T.I., Beshlej, I.V., Deryagina, V.P., Ryzhova, N.I., & Matistov, N.V. (2012). Izuchenie khimicheskogo sostava listev *schoenoprasum* L. i ingibiruiushchego deistviia ikh ekstraktov na opukholevyi rost u myshei [Study of the chemical composition of *A. schoenoprasum* L. leaves and the inhibitory effect of their extracts on tumor growth in mice]. *Khimiko-farmatsevticheskii zhurnal — Chemico-Pharmaceutical Journal*, 46(11), 40–43. DOI: <https://doi.org/10.30906/0023-1134-2012-46-11-30-33> [in Russian].
- 9 Matistov, N.V. (2013). Mikronutrienty dikorastushchikh i kultiviruemykh vidov roda *Allium* (*A. angulosum*, *A. strictum*, *A. schoenoprasum*) na evropeiskom Severo-Vostoke Rossii [Micronutrients of wild and cultivated species of the genus *Allium* (*A. angulosum*, *A. strictum*, *A. schoenoprasum*) in the European North-East of Russia]. *Extended abstract of candidate's thesis*. Novosibirsk [in Russian].
- 10 Jang, J.E., Baasanmunkh, S., Nyamgerel, N., Oh, S.Y., Song, J.H., Yusupov, Z., Tojibaev, K., & Choi, H.J. (2024). Flower morphology of *Allium* (Amaryllidaceae) and its systematic significance. *Plant Diversity*, 46(1), 3–27. DOI: 10.1016/j.pld.2023.06.009.
- 11 Ivanova, M.I., Baikov, A.A., Gins, E.M., Gins, V.K., Kashleva, A.I., Gins, M.S., & Smurova, N.V. (2024). Assessment of phytochemicals in *Allium* species: a systematic review. *Sabrao Journal of Breeding & Genetics*, 56(3). DOI: 10.54910/sabrao2024.56.3.13.
- 12 Teotia, D., Agrawal, A., Goyal, H., Jain, P., Singh, V., Verma, Y., Parveen, K., Bukhari, N.A., Chandra, A., & Malik, V. (2024). Pharmacophylogeny of genus *Allium* L. *Journal of King Saud University-Science*, 36(8), 103330. DOI: 10.1016/j.jksus.2024.103330.
- 13 Feknous, N., Boumendjel, M., & Leblab, F.Z. (2024). Updated Insights on the Antimicrobial Activities of *Allium* Genus (A Review). *Russian Journal of Bioorganic Chemistry*, 50(3), 806–823. DOI: 10.1134/S1068162024030257.
- 14 Iwar, K., Ochar, K., Seo, Y.A., Ha, B.K., & Kim, S.H. (2024). *Allium* as Potential Antioxidants and Anticancer Agents. *International Journal of Molecular Sciences*, 25(15), 8079. DOI: 10.3390/ijms25158079.
- 15 Friesen, N., Herden, T., Leweke, M., Grützmacher, L., Fragman-Sapir, O., Hurka, H., Blattner, F.R., & Fritsch, R.M. (2024). Dated phylogeny, phylogeography, and classification of *Allium* subgenus *Amerallium* (Amaryllidaceae) from the Old World, based on six DNA fragments. *Taxon*, 73(4), 971–991. DOI: <https://doi.org/10.1002/tax.13217>.
- 16 Vatandaşlar, Ö. & Koçyiğit, M. (2024). Comparative Seed Morphology of Eight *Allium* L. Species from Sections Codonoprasum and Scorodon. *Turkish Journal of Bioscience and Collections*, 8(2), 41–48. DOI: <https://doi.org/10.26650/tjbc.1525986>.
- 17 Baitulin, I.O. & Sitpaeva, G.T. (Eds.). (2014). *Krasnaia kniga Kazakhstana [Red Book of Kazakhstan]*. Vol. 2. Part 1. *Rasteniia — Plants*. (2d ed.). Astana: TOO «ArtPrintXXI» [in Russian].

- 18 Gemejjeva, N.G., Tokenova, A.M., & Frizen, N.V. (2021). Obzor sovremennogo sostoiianiia i perspektivy izucheniia kazakhstanskikh vidov roda *Allium* L. [Review of the current state and prospects of studying Kazakh species of the genus *Allium* L.]. *Problemy botaniki Yuzhnoi Sibiri i Mongolii — Problems of Botany of Southern Siberia and Mongolia*, 20, 1, 97–101. DOI: <https://doi.org/10.14258/pbssm.2021020> [in Russian].
- 19 Rachkovskaya, E.I., Safronova, I.N., & Volkova, E.A. (2003). Printsipy i osnovnye edinitzy raionirovaniia [Principles and basic units of zoning]. *Botanicheskaia geografiia Kazakhstana i Srednei Azii (v predelakh pustynnoi oblasti) — Botanical geography of Kazakhstan and Central Asia (within the desert region)*. Saint Petersburg [in Russian].
- 20 Korchagin, A.A. (1964). Vidovoi (floristicheskii) sostav rastitelnykh soobshchestv i metody ego izucheniia [Species (floristic) composition of plant communities and methods of its study]. *Polevaia geobotanika — Field geobotany*, 3, 39–60. Moscow-Leningrad [in Russian].
- 21 Ponyatovskaya, V.M. (1964). Uchet obilii i osobennosti razmeshcheniia vidov v estestvennykh rastitelnykh soobshchestvakh [Consideration of abundance and peculiarities of species placement in natural plant communities]. *Polevaia geobotanika — Field geobotany*, 3, 209–237. Moscow-Leningrad [in Russian].
- 22 Bykov, B.A. (1978). Geobotanika [Geobotany]. Alma-Ata: Nauka [in Russian].
- 23 Pavlov, N.V. (Ed.). (1956). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 1. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 24 Pavlov, N.V. (Ed.). (1958). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 2. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 25 Pavlov, N.V. (Ed.). (1960). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 3. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 26 Pavlov, N.V. (Ed.). (1961). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 4. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 27 Pavlov, N.V. (Ed.). (1961). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 5. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 28 Pavlov, N.V. (Ed.). (1963). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 6. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 29 Pavlov, N.V. (Ed.). (1964). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 7. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 30 Pavlov, N.V. (Ed.). (1965). Flora Kazakhstana [Flora of Kazakhstan]. (Vols. 1–9). Vol. 8. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 31 Pavlov, N.V. (Ed.). (1966). Flora Kazakhstana [Flora of Kazakhstan]. (Vol. 1–9). Vol. 9. Alma-Ata: Izdatelstvo Akademii nauk Kazakhskoi SSSR [in Russian].
- 32 Goloskokov, V.P. (Ed.). (1969). Illiustrirovannyi opredelitel rastenii Kazakhstana [Illustrated Identifier of Plants of Kazakhstan]. (Vols. 1–2). Vol. 1. Alma-Ata, Izdatelstvo “Nauka” Kazakhskoi SSR [in Russian].
- 33 Goloskokov, V.P. (Ed.). (1969). Illiustrirovannyi opredelitel rastenii Kazakhstana [Illustrated Identifier of Plants of Kazakhstan]. (Vols. 1–2). Vol. 2. Alma-Ata, Izdatelstvo “Nauka” Kazakhskoi SSR [in Russian].
- 34 (2007–2024). Plantarium: otkryti online atlas-opredelitel rastenii i lichainikov Rossii i sopredelnykh stran [Plantarium: an open online atlas-definition of plants and lichens of Russia and neighbouring countries]. Retrieved from <http://www.plantarium.ru/> [in Russian].
- 35 (2024). Plants of the World Online. POWO. Retrieved from <https://powo.science.kew.org>
- 36 Sokolov, P.D. (Ed.). (1994). Rastitelnye resursy Rossii i sopredelnykh gosudarstv: Tsvetkovye rasteniia, ikh khimicheskii sostav, ispolzovanie [Plant resources of Russia and neighboring countries. Flowering plants and their chemical composition, use]. *Semeistva: Butomaceae — Typhaceae*. Saint Petersburg: Nauka [in Russian].
- 37 Budancev, A.L. (Ed.). (1996). Rastitelnye resursy Rossii i sopredelnykh gosudarstv [Plant resources of Russia and neighbouring countries]. *Chast 1: Semeistva Lycopodiaceae — Ephedraceae; Chast 2: Dopilnenie k 1-7 tomam — Part 1: Lycopodiaceae — Ephedraceae families; Part 2: Supplements to Volumes 1–7*. Saint Petersburg: Mir i semia–95 [in Russian].
- 38 Budancev, A.L. (Ed.). (2008). Rastitelnye resursy Rossii: Dikorastushchie tsvetkovye rasteniia, ikh komponentnyi sostav i biologicheskaiia aktivnost [Plant Resources of Russia: Wild Flowering Plants, Their Component Composition and Biological Activity]. *Tom 1. Semeistva: Magnoliaceae — Juglandaceae, Ulmaceae, Moraceae, Cannabaceae, Urticaceae — Volume 1. Families of Magnoliaceae — Juglandaceae, Ulmaceae, Moraceae, Cannabaceae, Urticaceae*. Saint Petersburg, Moscow: Tovarishchestvo nauchnykh izdaniy KMK [in Russian].
- 39 Budancev, A.L., & Lesiovskaya, E.E. (2001). Dikorastushchie poleznye rasteniia Rossii [Wild useful plants of Russia]. Saint Petersburg: Izdatelstvo SPKhFA [in Russian].
- 40 Grudzinskaya, L.M., Gemedzhieva, N.G., Nelina, N.V., & Karzhaubekova, Zh.Zh. (2014). *Annotirovannyi spisok lekarstvennykh rastenii Kazakhstana: Spravochnoe izdanie [Annotated list of medicinal plants of Kazakhstan: Reference edition]*. Almaty [in Russian].
- 41 Vvedensky, A.I. (1971). Rod *Allium* L. [Genus *Allium* L.]. *Opredelitel rastenii Srednei Azii — Identifier of plants of Central Asia*, 2, 39–89. Tashkent, Izdatelstvo FAN Uzbekskoi SSR [in Russian].
- 42 Aralbay, N.K., Mukhtybaeva, A.A., & Shormanova, A.A. (2007). *Krasnaia kniga Zhambylskoi oblasti [Red Data Book of Zhambyl region]*. Almaty [in Russian].

- 43 Lazkov, G.A., & Sultanova, B.A. (2014). *Kadastr flory Kyrgyzstana. Sosudistye rasteniia [Inventory of the flora of Kyrgyzstan. Vascular plants]*. Bishkek [in Russian].
- 44 Turdumatova, N.K. (2022). Rod luk (*Allium* L.) v Kyrgyzstane (voprosy sistematiki, geografiia i perspektivy ispolzovaniia): [Genus *Allium* L. in Kyrgyzstan (questions of systematics, geography, and prospects of use)]. *Extended abstract of candidate's thesis*. Bishkek [in Russian].
- 45 Friesen, N., Grützmacher, L., Skaptsov, M., Vesselova, P., Dorofeyev, V., Luferov, A.N., Turdumatova, N., Lazkov, G., Smirnov, S.V., Shmakov, A.I., & Hurka, H. (2022). *Allium pallasii* and *A. caricifolium* — surprisingly diverse old steppe species, showing a clear geographical barrier in the area of Lake Zaysan. *Plants*, 11(11), 1465. doi.org/10.3390/plants11111465.
- 46 Nelina, N.V., Kudabayeva, G.M., Mukhtubayeva, S.K., & Bilibayeva, B.K. (2016). K vidovomu sostavu flory ushchelii Merke i Karakystak zapadnoi chasti Kirgizskogo Alatau [The species composition of the flora of the Merke and Karakystak gorges of the western part of the Kyrgyz Alatau]. *Problemy botaniki Yuzhnoi Sibiri i Mongolii: materialy XV Mezhdunarodnoi nauchno-prakticheskoi konferentsii — Problems of botany of South Siberia and Mongolia: materials of the XV International scientific and practical conference* (pp. 392–398) [in Russian].

Information about the authors

Gemejiyeva Nadezhda Gennadievna — Doctor of Biological Sciences (D.Sc.), Professor, Head of Plant Resource Laboratory, RSE REM “Institute of Botany and Phytointroduction” FWC MENR RK, Timiryazev street, 36 “D”, 050040, Almaty, Kazakhstan; e-mail: ngemed58@mail.ru; ORCID: 0000-0002-7317-2685

Abdildanov Daulet Sharipkhanovich — Phd doctoral student, Department of Biodiversity and Bioresources, Al-Farabi Kazakh National University, Timiryazeva 71, 050040, Almaty, Kazakhstan; Junior Researcher at the Laboratory of Higher Plant Flora, RSE REM “Institute of Botany and Phytointroduction” FWC MENR RK, Timiryazev street, 36 “D”, 050040, Almaty, Kazakhstan; e-mail: abdildanov00@mail.ru; ORCID: 0000-0002-0681-5468

Kudabayeva Gulmira Mauletovna — Candidate of Biological Sciences (C. Sc), Leading Researcher, Laboratory of Higher Plant Flora, RSE REM “Institute of Botany and Phytointroduction” FWC MENR RK, Timiryazev street, 36 “D”, 050040, Almaty, Kazakhstan; e-mail: kgm_anita@mail.ru; ORCID: 0000-0001-7315-4033

Vesselova Polina Vasilyevna — Candidate of Biological Sciences (C. Sc), Head of the Laboratory of Flora of Higher plants RSE REM “Institute of Botany and Phytointroduction” FWC MENR RK, Timiryazev street, 36 “D”, 050040, Almaty, Kazakhstan; e-mail: pol_ves@mail.ru; ORCID: 0000-0002-3903-6276

Nelina Natalia Vasilyevna — Candidate of Biological Sciences (C. Sc), Senior Research Fellow, Laboratory of Higher Plant Flora, RSE REM “Institute of Botany and Phytointroduction” FWC MENR RK, Timiryazev street, 36 “D”, 050040, Almaty, Kazakhstan; e-mail: nelina.natalia@mail.ru; ORCID: 0000-0001-9064-881X

Friesen Nikolay Valterovich — Doctor of Biological Sciences (D.Sc.), Professor, Botanical Garden, University of Osnabrück, Albrechtstrasse 29, 49076, Osnabrück, Germany; e-mail: nfriesen@uni-osnabrueck.de. ORCID: 0000-0003-3547-3257